


EXODUS

# EXODUS

AUTOMATIC EXOSOME  
ISOLATION SYSTEM






---

## Automatic System for Exosomes Isolation

---



### EXODUS

EXODUS is an automatic, label-free, and highly efficient exosome isolation system. With EXODUS, you can easily and quickly isolate high-quality, intact exosomes with excellent yield and purity from a variety of bio-fluids and sample volumes.





Experience the efficiency of EXODUS for yourself and take your research to the next level.





## Isolation Principles

EXODUS has been developed using a dual-membrane nanofiltration system that integrates periodic negative pressure oscillation (NPO) and double-coupled ultrasonic harmonic oscillations (HO).



Nature Methods, 2021, 18(2):212-218.


EXODUS can rapidly remove free nucleic acid and protein impurities from the sample, resulting in the efficient purification and enrichment of exosomes. The exosomes are precisely intercepted by nanoporous membrane, allowing for a highly targeted isolation process.

EXODUS has great potential to revolutionize exosome isolation and drive new discoveries in biomedical research and translation.

# Automatic

EXODUS is designed to automatically isolate high yield and purity exosomes from different biofluid sample volumes.

## Step 1



### Sample loading



Plasma



Urine



Saliva



Tears




Aqueous  
humor




Synovial  
fluid

### Automatic isolation



## Step 2




## Step 3



### Easy exosome collection

### TEM





# KEY FEATURES

**EXODUS**  
Key  
Features>>





### Rapid isolation

Maximum isolation speed: 200 mL/h

---



### High purity and high yield

Purity ~ 99 %; Yield ~ 90 %

---



### Wide application

Sample types	Sample volumes	Sample types	Sample volumes
Urine	1 - 250 mL	Plasma	0.01 - 2 mL
Plant		Saliva	0.5 - 10 mL
Cell culture medium		Tears	0.005 - 1 mL
Cell-derived vesicle		Aqueous humor	0.005 - 1 mL
Bacterial culture medium		Cerebrospinal fluid	0.5 - 25 mL

---




### Label-free

Only need PBS buffer




Rapid isolation



Nutture Methods, 2021, 18(2):212-218.



High purity and high yield



Nutture Methods, 2021, 18(2):212-218.



Wide application

Other small amount sample types



Plasma



Tears



Saliva



Cerebrospinal fluid



Aqueous humor




Synovial fluid



Label-free

TEM image of exosome





**EXODUS**  
Application >>

**APPLICATION**



---

## Various Sample Types

---



Plasma



Urine



Saliva



Cerebrospinal  
fluid



Tears



Aqueous  
humor



Synovial  
fluid



Tissue



Cell culture  
medium



Bacterial  
culture medium



Cell-derived  
vesicle



Plant

...

---

## Applications


---

- Early diagnosis
- Drug delivery
- Exosome therapeutics
- Regenerative medicine

## 1 Urine EV Isolation with EXODUS


**Figure 1. Characterization of exosomes by transmission electron microscopy (TEM).**

TEM of exosomes harvested from urine by EXODUS, showing the characteristic cup and plate shape of exosomes. Scale bar = 600 nm.



**Figure 2. Method comparison for exosome isolation.**

Western blot analysis of exosomal markers (Alix, CD63, TSG101, and CD81) and impurity Uromodulin (UMOD) of exosomes isolated using methods of EXODUS, ultracentrifugation (UC), PEG precipitation, Membrane affinity (MA), Size exclusion chromatography (SEC), and phosphatidylserine (PS) affinity, respectively. Compared to other methods, exosome isolated by EXODUS shows higher relative yield and purity in WB gel graph (left) and bar graph of normalized band intensity (right).




Nature Methods, 2021, 18(2):212-218.

## 2 Cell culture medium EV Isolation with EXODUS


**Figure 1. Characterization of exosomes by TEM.**

Electron micrographs of exosomes isolated from umbilical cord mesenchymal stem cell supernatants using EXODUS H (left) and UC (Right). TEM of exosomes obtained from EXODUS shows such typical cupped structure, more exosomal vesicular particles, and less particle aggregation compared to that from UC. Scale bar = 600 nm.



**Figure 2. Scratched wound assay for the effects evaluation of exosomes on migration capacity of cells.**

Cells began to exhibit significantly migration 12h after treatment with exosomes isolated either from EXODUS or EXO-kit compared to that from UC and the control group. After 24h, enhanced cell migration was observed from the group with the addition of EXODUS-isolated exosome. Scale bar = 40  $\mu$ m.




Electrophoresis, 2024, 0173-0835.

### 3 Plasma EV Isolation with EXODUS

**Figure 1. Characterization of exosomes by TEM.**


TEM analysis of exosomes isolated from the plasma of patients with esophageal squamous cell carcinoma using the EXODUS reveals their distinctive cup-shaped and plate-like morphology. Scale bar = 400 nm.



Technol Cancer Res Treat, 2024, 1533-0338.

**Figure 2. The yield and purity analysis of sEVs isolated by methods of EXODUS, UC, and PEG, respectively.**

sEVs isolated using EXODUS or PEG from 20  $\mu\text{L}$  of plasma (approximately  $2 \times 10^{10}$  particles/mL) showed over a 50-fold increase compared to those isolated using UC. The purity of sEVs isolated using EXODUS was more than three times higher than that achieved with UC or PEG.




Int J Nanomedicine, 2024, 19:1351-1362.

### 4 Trace sample EV Isolation with EXODUS

**Figure 1. Characterization of tears EVs and aqueous humor EVs by TEM.**

TEM images of exosomes derived from two trace samples of 50  $\mu\text{L}$  tear fluid (left, scale bar = 100 nm) and 150  $\mu\text{L}$  aqueous humor (right, scale bar = 400 nm) using the EXODUS, reveals their characteristic cup-shaped and plate-like morphological features.




ACS Nano, 2022, 16(8): e11720.

**Figure 2. The yield and purity analysis of tear exosome isolated by methods of EXODUS, UC, SEC, and PEG.**

**A.** Total particles of exosomes isolated using EXODUS or PEG from 50  $\mu\text{L}$  of tear (approximately  $2 \times 10^{10}$  particles/mL) showed over a 20-fold and 2-fold increase compared to those isolated from UC and SEC, respectively.

**B.** Equal-protein-mass (3  $\mu\text{g}$ ) western blot analysis of the exosomal markers (Alix, CD63, CD9, and CD81) and negative marker (Calnexin) of tear exosomes prepared by iTEARS EXODUS and other methods.




ACS Nano, 2022, 16(8): e11720.



## 5 Bacterial EV Isolation using with EXODUS


**Figure 1. Characterization of bacterial EVs by TEM.**

TEM images of EVs derived from *Helicobacter pylori*, *Escherichia coli*, *Mycobacterium avium*, and intestinal microbiota isolated by EXODUS H 600. The EXODUS H 600 is capable of isolating EVs from a variety of bacterial sources, including both Gram-positive and Gram-negative bacteria.



**Figure 2. Comparison of yield and purity by using EXODUS and UC.**


EXODUS achieves higher particle yield and EV purity compared to UC of *E. coli* EVs.



## 6 Virus and virus-like particles Isolation with EXODUS


**Figure 1. TEM characterization of virus and virus-like particles.**

Using EXODUS, high-resolution TEM images were obtained to highlight typical morphology of bacteriophages, phage protein self-assembly particles and Rift Valley fever virus. Gentle isolation of EXODUS is able to preserve the viral or particle integrity, which is crucial for accurate structural analysis. Scale bar = 100 nm.



**Figure 2. Lentiviral titer recovery rate results**

After purification with the EXODUS, the titer of lentiviral infection increased from  $9.86 \times 10^7$  to  $1.56 \times 10^9$ , with a recovery rate of 63.29%, and the physical titer increased from  $2.5 \times 10^{10}$  to  $3.95 \times 10^{11}$ , with a recovery rate of 63.0%.



# Publications

NO.	Title	Journal	Five-year average IF
01	Exosome detection via the ultrafast-isolation system: EXODUS	Nature Methods	45.6
02	UPCARE:Urinary Extracellular Vesicles-Derived Prostate Cancer Assessment for Risk Evaluation	Journal of Extracellular Vesicles	19.6
03	Discovering the Secret of Diseases by Integrated Tear Exosomes Analysis via Rapid-isolation System: iTEARS	ACS Nano	16.2
04	Robust Acute Pancreatitis Identification and Diagnosis: RAPIDx	ACS Nano	16.2
05	Interaction network of extracellular vesicles building universal analysis via eye tears: iNEBULA	Science Advances	13.7
06	Sensitive small extracellular vesicles associated circRNAs analysis combined with machine learning for precision identification of gastric cancer	Chemical Engineering Journal	13.2
07	Quantitative metabolic analysis of plasma extracellular vesicles for the diagnosis of severe acute pancreatitis	Journal of Nanobiotechnology	11.4
08	Lipidomic identification of urinary extracellular vesicles for non-alcoholic steatohepatitis diagnosis	Journal of Nanobiotechnology	11.4
09	Metabolomic investigation of urinary extracellular vesicles for early detection and screening of lung cancer	Journal of Nanobiotechnology	11.4
10	The genetic source tracking of human urinary exosomes	PNAS	10.8
11	Identification and detection of plasma extracellular vesicles-derived biomarkers for esophageal squamous cell carcinoma diagnosis	Biosensors & Bioelectronics	9.9
12	Audible Acoustic Wave Promotes EV Formation and Secretion from Adherent Cancer Cells via Mechanical Stimulation	ACS Applied Materials & Interfaces	8.7
13	Identification of circulating extracellular vesicle long RNAs as diagnostic biomarkers for patients with severe acute pancreatitis	Clinical and Translational Medicine	8.0
14	Prediction of Response to Chemoradiotherapy by Dynamic Changes of Circulating Exosome Levels in Patients with Esophageal Squamous Cell Carcinoma	International Journal of Nanomedicine	7.5
15	Investigating the proliferative inhibition of HepG2 cells by exosome-like nanovesicles derived from Centella asiatica extract through metabolomics	Biomedicine & Pharmacotherapy	6.8
16	Gut Subdoligranulum variabile ameliorates rheumatoid arthritis by promoting TSG-6 synthesis from joint cells	Frontiers in Immunology	6.8
17	Metabolomic analysis of exosomal-markers in esophageal squamous cell carcinoma	Nanoscale	6.1
18	Metabolic signatures of tear extracellular vesicles caused by herpes simplex keratitis	Ocular Surface	5.9
19	Sensitive electrochemical biosensor for rapid detection of sEV-miRNA based turbo-like localized catalytic hairpin assembly	Analytica Chimica Acta	5.5
20	Isolation of Exosome Nanoparticles from Human Cerebrospinal Fluid for Proteomic Analysis	ACS Applied Nano Materials	5.4
21	Human umbilical cord mesenchymal stem cells inhibit liver fibrosis via the microRNA-148a-5p/SLIT3 axis	International Immunopharmacology	5.0
22	Patient-derived induced pluripotent stem cells with a MERTK mutation exhibit cell junction abnormalities and aberrant cellular differentiation potential	World Journal of Stem Cells	4.2
23	Assessing Alzheimer's disease via plasma extracellular vesicle-derived mRNA	Alzheimer's & Dementia: Diagnosis, Assessment & Disease Monitoring	4.0
24	Efficient preparation of high-purity and intact mesenchymal stem cell-derived extracellular vesicles	Analytical and Bioanalytical Chemistry	3.8
25	Plasma-derived Exosomal miR-25-3p and miR-23b-3p as Predictors of Response to Chemoradiotherapy in Esophageal Squamous Cell Carcinoma	Technology in Cancer Research & Treatment	2.8
26	Comparative investigation of exosome extraction from rat bone marrow mesenchymal stem cells using three different methodologies	Electrophoresis	2.8
27	Isolation of small extracellular vesicles from a drop of plasma via EXODUS and their fingerprint proteomics profiling by MALDI-TOF MS	Biosensors & Bioelectronics:X	/
28	Application of EXODUS system combined with allosteric DNA nanoswitches in the detection of miR-107 among plasma exosomes of Parkinson's disease patients	Chinese Journal of Preventive Medicine	/

# EXODUS System Specification >>

# SPECIFICATION



Model	EXODUS H-300	EXODUS H-600
Isolation principles	Combination of the negative pressure oscillations (NPO) and double coupled harmonic oscillations (HO) on nanoporous membrane	
Sample types	Plasma, urine, saliva, cerebrospinal fluid, tears, aqueous humor, synovial fluid, tissue, cell culture medium, bacterial culture medium, cell-derived vesicle, plant, ect.	
Isolation device size	S/M	S/M/L
Temperature of sample reservoir	2 - 8 °C	
Sample volumes	10 µL - 50 mL	10 µL - 250 mL
Processing speed	Max speed 50 mL/h	Max speed 200 mL/h
Isolation data saving	2000	20000
Exosome recovery volumes	100 - 400 µL	100 - 1000 µL
Ultraviolet sterilization	Internal UV lamp, turn off automatically after 30 min	
Display	10.4 inch touch screen, real time display with sample type, time, processing information ect. Supporting the operation without computer	
Dimension	535 x 510 x 475 mm (H x W x D)	
Net weight	40 kg (88 lbs)	
System interfaces	4 USB ports, 1 network port, 1 serial port	



# EXODUS

Product specifications may change without notice,  
based on the latest technical data and test results.

✉ [service@exodus-bio.com](mailto:service@exodus-bio.com)

📍 160 E Tasman Dr., San Jose, CA 95134, United States